

§ 520.23

520.2134 Spinosad and milbemycin.
 520.2150 Stanazolol.
 520.2158 Streptomycin.
 520.2170 Sulfabromomethazine.
 520.2184 Sulfachloropyrazine.
 520.2200 Sulfachlorpyridazine.
 520.2215 Sulfadiazine/pyrimethamine sus-
 pension.
 520.2218 Sulfamerazine, sulfamethazine, and
 sulfaquinoxaline powder.
 520.2220 Sulfadimethoxine oral dosage
 forms.
 520.2220a Sulfadimethoxine solution and
 soluble powder.
 520.2220b Sulfadimethoxine suspension.
 520.2220c Sulfadimethoxine tablet.
 520.2220d Sulfadimethoxine bolus.
 520.2220e Sulfadimethoxine extended-release
 bolus.
 520.2220f Sulfadimethoxine and ormetoprim
 tablet.
 520.2240 Sulfaethoxyypyridazine.
 520.2240a Sulfaethoxyypyridazine solution.
 520.2240b Sulfaethoxyypyridazine tablets.
 520.2260 Sulfamethazine oral dosage forms.
 520.2260a Sulfamethazine oblet, tablet, and
 bolus.
 520.2260b Sulfamethazine sustained-release
 boluses.
 520.2260c Sulfamethazine sustained-release
 tablets.
 520.2261 Sulfamethazine sodium oral dosage
 forms.
 520.2261a Sulfamethazine solution.
 520.2261b Sulfamethazine powder.
 520.2280 Sulfamethizole and methenamine.
 520.2325 Sulfaquinoxaline oral dosage forms.
 520.2325a Sulfaquinoxaline powder and solu-
 tion.
 520.2325b Sulfaquinoxaline drench.
 520.2330 Sulfisoxazole tablets.
 520.2340 Tepoxalin.
 520.2345 Tetracycline.
 520.2345a Tetracycline capsules.
 520.2345b Tetracycline tablets.
 520.2345c Tetracycline boluses.
 520.2345d Tetracycline powder.
 520.2345e Tetracycline solution.
 520.2345f Tetracycline phosphate complex
 and sodium novobiocin capsules.
 520.2345g Tetracycline hydrochloride and so-
 dium novobiocin tablets.
 520.2345h Tetracycline hydrochloride, so-
 dium novobiocin, and prednisolone tab-
 lets.
 520.2362 Thenium closylate.
 520.2380 Thiabendazole oral dosage forms.
 520.2380a Thiabendazole top dressing and
 mineral protein block.
 520.2380b Thiabendazole drench or paste.
 520.2380c Thiabendazole bolus.
 520.2380d Thiabendazole and piperazine cit-
 rate.
 520.2380e Thiabendazole and trichlorfon.
 520.2380f Thiabendazole and piperazine phos-
 phate.
 520.2455 Tiamulin.

21 CFR Ch. I (4–1–15 Edition)

520.2471 Tilmicosin.
 520.2473 Tioxidazole oral dosage forms.
 520.2473a Tioxidazole granules.
 520.2473b Tioxidazole paste.
 520.2475 Toceranib.
 520.2481 Triamcinolone acetonide tablets.
 520.2482 Triamcinolone acetonide oral pow-
 der.
 520.2483 Triamcinolone.
 520.2520 Trichlorfon oral dosage forms.
 520.2520a Trichlorfon and atropine.
 520.2520b Trichlorofon boluses.
 520.2520c Trichlorofon granules.
 520.2520d Trichlorfon, phenothiazine, and pi-
 perazine dihydrochloride powder.
 520.2582 Triflupromazine.
 520.2598 Trilostane.
 520.2604 Trimeprazine and prednisolone tab-
 lets.
 520.2605 Trimeprazine and prednisolone cap-
 sules.
 520.2610 Trimethoprim and sulfadiazine tab-
 lets.
 520.2611 Trimethoprim and sulfadiazine
 paste.
 520.2612 Trimethoprim and sulfadiazine sus-
 pension.
 520.2613 Trimethoprim and sulfadiazine
 powder.
 520.2640 Tylosin.
 520.2645 Tylvalosin.

AUTHORITY: 21 U.S.C. 360b.

SOURCE: 40 FR 13838, Mar. 27, 1975, unless
 otherwise noted.

§ 520.23 Acepromazine.

(a) *Specifications*. Each tablet con-
 tains 5, 10, or 25 milligrams (mg)
 acepromazine maleate.

(b) *Sponsors*. See No. 000010 in
 § 510.600(c) of this chapter.

(c) *Conditions of use*(1) *Dogs*—(i)
Amount. 0.25 to 1.0 mg per pound (lb)
 body weight orally.

(ii) *Indications for use*. As an aid in
 tranquilization and as a preanesthetic
 agent.

(iii) *Limitations*. Federal law restricts
 this drug to use by or on the order of a
 licensed veterinarian.

(2) *Cats*—(i) *Amount*. 0.5 to 1.0 mg/lb
 body weight orally.

(ii) *Indications for use*. As a tranquil-
 izer.

(iii) *Limitations*. Federal law restricts
 this drug to use by or on the order of a
 licensed veterinarian.

[75 FR 10165, Mar. 5, 2010]

§ 520.28 Acetazolamide.

(a) *Specifications*. A powder con-
 taining acetazolamide sodium, USP

Food and Drug Administration, HHS

§ 520.38b

equivalent to 25 percent acetazolamide activity.

(b) *Sponsor*. See No. 054771 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs*—(1) *Amount*. Administer orally at a dosage of 5 to 15 milligrams per pound of body weight daily.

(2) *Indications for use*. As an aid in the treatment of mild congestive heart failure and for rapid reduction of intraocular pressure.

(3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[79 FR 28816, May 20, 2014]

§ 520.38 Albendazole oral dosage forms.

§ 520.38a Albendazole suspension.

(a) *Specifications*. Each milliliter of suspension contains 45.5 milligrams (mg) (4.55 percent) or 113.6 mg (11.36 percent) albendazole.

(b) *Sponsor*. See No. 054771 in § 510.600 of this chapter.

(c) *Related tolerances*. See § 556.34 of this chapter.

(d) *Special considerations*. See § 500.25 of this chapter.

(e) *Conditions of use*—(1) *Cattle*. Administer 11.36 percent suspension:

(i) *Amount*. 4.54 mg/pound (lb) body weight (10 mg/kilogram (kg)) as a single oral dose using dosing gun or dosing syringe.

(ii) *Indications for use*. For removal and control of adult liver flukes (*Fasciola hepatica*); heads and segments of tapeworms (*Moniezia benedeni* and *M. expansa*); adult and 4th stage larvae of stomach worms (brown stomach worms including 4th stage inhibited larvae (*Ostertagia ostertagi*), barberpole worm (*Haemonchus contortus* and *H. placei*), small stomach worm (*Trichostrongylus axei*)); adult and 4th stage larvae of intestinal worms (thread-necked intestinal worm (*Nematodirus spathiger* and *N. helvetianus*), small intestinal worm (*Cooperia punctata* and *C. oncophora*)); adult stages of intestinal worms (hookworm (*Bunostomum phlebotomum*), bankrupt worm (*Trichostrongylus colubriformis*), nodular worm (*Oesophagostomum radiatum*)); adult and 4th stage larvae of lungworms (*Dictyocaulus viviparus*).

(iii) *Limitations*. Do not slaughter within 27 days of last treatment. Do not use in female dairy cattle of breeding age; Do not administer to female cattle during first 45 days of pregnancy or for 45 days after removal of bulls.

(2) *Sheep*. Administer 4.45 or 11.36 percent suspension:

(i) *Amount*. 3.4 mg/lb body weight (7.5 mg/kg) as a single oral dose using dosing gun or dosing syringe.

(ii) *Indications for use*. For removal and control of adult liver flukes (*Fasciola hepatica* and *Fascioloides magna*); heads and segments of common tapeworms (*Moniezia expansa*) and fringed tapeworm (*Thysanosoma actinioides*); adult and fourth stage larvae of stomach worms (brown stomach worm (*Ostertagia circumcincta* and *Marshallagia marshalli*), barberpole worm (*Haemonchus contortus*), small stomach worm (*Trichostrongylus axei*)); adult and fourth stage larvae of intestinal worms (thread-necked intestinal worm (*Nematodirus spathiger* and *N. filicollis*), Cooper's worm (*Cooperia oncophora*), bankrupt worm (*Trichostrongylus colubriformis*), nodular worm (*Oesophagostomum columbianum*), and large-mouth bowel worm (*Chabertia ovina*)); adult and larval stages of lungworms (*Dictyocaulus filaria*).

(iii) *Limitations*. Do not slaughter within 7 days of last treatment. Do not administer to ewes during first 30 days of pregnancy or for 30 days after removal of rams.

(3) *Goats*. Administer 11.36 percent suspension:

(i) *Amount*. 4.54 mg/lb body weight (10 mg/kg) as a single oral dose using dosing gun or dosing syringe.

(ii) *Indications for use*. For the treatment of adult liver flukes (*Fasciola hepatica*) in nonlactating goats.

(iii) *Limitations*. Do not slaughter within 7 days of last treatment. Do not administer to does during the first 30 days of pregnancy or for 30 days after removal of bucks.

[73 FR 11027, Feb. 29, 2008. Redesignated at 78 FR 66264, Nov. 5, 2013, as amended at 79 FR 28816, May 20, 2014]

§ 520.38b Albendazole paste.

(a) *Specifications*. The product contains 30 percent albendazole.